



## **MMS PROTOCOL 2000 –**

This Protocol is for all cancers and other life threatening diseases. It basically is taking 3 activated drops each hour without becoming any sicker, or feeling worse, or developing diarrhea. It must be done for 8-10 hours a day, for 3 weeks or until well. It is best to start off with only one activated drop an hour and quickly build up to 3, or as many as you can tolerate, again, without feeling worse.

**PLEASE NOTE:** There is no benefit in making yourself sicker than you already are. This uses up energy and makes the recovery go slower. If you notice any of these symptoms; nausea, pain, or diarrhea, reduce the number of activated drops until the symptoms disappear. Try not to stop taking MMS, just take fewer drops.

After 3 days of successfully taking MMS1 drops hourly for ten hours a day, you must add MMS2 Capsules. MMS2 capsules are Calcium Hypochlorite. When dissolved in water it produces Hypochlorous Acid.

(\*\* With MMS2 capsules you may want to start slowly, with ¼ capsule for the first day, and ½ capsule for the second day, and then a full capsule thereafter.)

When reaching the full dosing schedule, it will look something like this :

**Sample time schedule for Protocol 2000, once you have added MMS2 into your dosing:**

### **Protocol 2000**

#### **MMS2 Time Schedule**

##### **Time Dose**

**9:00 AM MMS1 dose**

**10:00 AM MMS1 dose**

**10:30 AM MMS2 dose**

**11:00 AM MMS1 dose**

**12:00 PM MMS1 dose**

**12:30 PM MMS2 dose**

**1:00 PM MMS1 dose**

**2:00 PM MMS1 dose**

**2:30 PM MMS2 dose**

**3:00 PM MMS1 dose**

**4:00 PM MMS1 dose**

**4:30 PM MMS2 dose**

**5:00 PM MMS1 dose**

**6:00 PM MMS1 dose**

**6:30 PM MMS2 dose**

**Note:** *While working up to the correct size capsules of MMS2, which is either a full #1 size capsule, or a 3/4 full #0 size capsule (**never go beyond these amounts**), keep your MMS1 doses constant. In other words, do not be working on increasing your drops of MMS1, while you are working up to your proper dose of MMS2, because if you get nauseous you will not be able to determine which of the two might be causing you to feel sick. Once you have reached the suggested amount of MMS2, then you can begin increasing your drops of MMS1 once more.*

*Remember, at any time, whether you are increasing your amounts of MMS1, or MMS2, if at any time you feel nauseous or sick from the increase, decrease the amount by at least 1/2 and build back up slowly.*

**\* Remember, Never take a dose containing DMSO and an MMS2 capsule at the same time! See page 13 of MMS Health Recovery Guide for the full warning on this.**

----

LEGAL DISCLAIMER: The information contained within emails or on our website are for informational purpose only. Genesis 2 Church Sacraments, services or technologies listed on this website are not medicines. They are not intended to diagnose, treat, cure or prescribe. None of our services, information or products are intended to replace the care or advise of a trained medical physician. You are solely responsible for how you use them.